

# HELM Nicosulfuron 75

Water Dispersible Granule Herbicide for  
Use on Corn

**ACTIVE INGREDIENT:**

Nicosulfuron 2-[[[4,6-dimethoxypyrimidin-2-yl]aminocarbonyl]aminosulfonyl] N,N-dimethyl-3-pyridinecarboxamide ..... 75%

**INERT INGREDIENTS:** ..... 25%

**TOTAL** ..... 100%

**STOP - Read the label before use**

**KEEP OUT OF REACH  
OF CHILDREN  
CAUTION**

EPA Reg. No. 74530-26 EPA Est. No. 083390-CHN-001

**Net Contents: 10 ou.**

**Distributed by:**

Helm Agro US, Inc.  
8295 Tournament Dr. Suite #310  
Memphis, TN 38125

Manufactured for:

**Sharda USA LLC**

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

**FIRST AID**

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **Have a product container or label with you when calling a poison control center or doctor, or going for treatment. Note To Physician: No specific antidote is available. Treat the patient symptomatically.**

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff.

**DIRECTIONS FOR USE**

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

**STORAGE AND DISPOSAL**

Do not contaminate water, other pesticides, fertilizer, food or feed in storage. **Pesticide Storage:** Store product in original container only. Store in a cool, dry place. **Pesticide Disposal:** Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** For Plastic Containers: Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. For Fiber Sacks: Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Dispose of bags at an approved waste disposal facility, in accordance with Federal, state and local regulations. For Fiber Drums With Liners: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. For Bags Containing Water Soluble Packets: Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. For Metal Containers (non aerosol): Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

#### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical resistant to this product are listed below. If you want more options follow the instructions for Category A on the EPA chemical resistance category selection chart.

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber), all > 14 mils.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

##### Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff.

## DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical resistant gloves Category A (such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber), all > 14 mils.
- Shoes plus socks.

HELM Nicosulfuron 75 should be used only in accordance with recommendations on this label or in supplemental Helm Agro US, INC. publications. Helm Agro US, INC. will not be responsible for losses or damage resulting from use of this product in any manner not specifically recommended by the guidelines provided on this label.

### **GENERAL INFORMATION**

- HELM Nicosulfuron 75 should be used at a rate of 1/3 – 1 1/3 oz. per acre for selective postemergence grass weed control in field corn grown for seed or grain, pop-corn and sweet corn.
- Do not make more than two (2) applications of HELM Nicosulfuron 75 per crop season
- The combined dosage of sequential applications of HELM Nicosulfuron 75 cannot exceed 1 1/3 oz. per acre.

### **WHEN TO APPLY:**

#### **Normal Planned Use**

HELM Nicosulfuron 75 may be used on field corn, high lysine, waxy, white or other food grade corn hybrids.

HELM Nicosulfuron 75 may be broadcast to corn up to 20" tall (free standing) or that is exhibiting up to and including 6 leaf collars (V6), whichever is more restrictive.

For best results, apply HELM Nicosulfuron 75 when applications are made early postemergence when corn and weeds are small. For best overall performance, apply HELM Nicosulfuron 75 to corn that is less than 12" tall.

#### **Timing to Weeds**

Apply HELM NICOSULFURON 75 when grasses are young and actively growing, but before they exceed the sizes indicated in Table 1. Treat heavy infestations of weeds before they become too competitive with the crop, especially where soil moisture and/or fertility are limited. HELM NICOSULFURON 75 provides weed control via foliar absorption. HELM NICOSULFURON 75 only controls those weeds that have emerged. For

later-emerging weeds, a second application or a timely cultivation is required. Applications made to weeds larger than the size indicated on this label or to weeds under stress may result in unsatisfactory control. Refer to LATE OR RESCUE APPLICATIONS.

### **LATE OR RESCUE APPLICATIONS**

HELM NICOSULFURON 75 may be applied to field corn as a rescue treatment for the control of escaped grasses, or as a directed postemergence application on corn that is taller than 20" or which has more than 6 collars (which ever occurs first). For corn 20" to 36" tall, apply HELM NICOSULFURON 75 with drop nozzles only and avoid spraying into the whorl of cornstalks.

Do not apply to corn that is taller than 36" or that exhibits 10 or more collars (V10), whichever is most restrictive.

Applications made to weeds larger than those listed on this label may vary from complete control to suppression. Level of control will depend on the weed species, stage of growth, and environmental conditions.

Due to the unplanned nature of rescue applications, choices must be made between the risks that arise from applications made beyond the proper time for HELM NICOSULFURON 75 use, and the effects of season long grass competition and/or harvest complications. These choices must balance risks from improperly timed HELM NICOSULFURON 75 use that include, but are not limited to:

- **Yield loss due to competition:** Research indicates competition from foxtail exceeding 4 inches in height may reduce corn yields. Applications to foxtail and other annual grasses that exceed the sizes stated on the label increases the risk of yield losses due to prolonged competition with the crop even though control may be acceptable.
- **Incomplete control of grasses beyond labeled size:** Applications to grasses that exceed the labeled sizes can result in reduced control. This incomplete control may reduce corn yield.

- **Incomplete grass control due to herbicide stress:** Grasses under stress from previous herbicide applications may not be actively growing and susceptible to HELM Nicosulfuron 75. This stress may reduce grass control in “rescue” situations.
- **Ear malformation:** Applications of HELM Nicosulfuron 75 on corn that has 7 to 10 collars (V7 to V10) increases the potential for ear malformation (pinching). This risk may be greatly reduced, but not eliminated, by using drop nozzles properly adjusted so as to not apply HELM Nicosulfuron 75 into the corn whorl.

#### **RATE**

Optimum control of the weeds listed can be achieved with 2/3 oz. of HELM Nicosulfuron 75. Weeds that exceed the listed weed sizes by up to 50% may be partially controlled with rates between 2/3 and 1 1/3 oz. of HELM Nicosulfuron 75 per acre.

HELM Nicosulfuron 75 may be applied at 1/3 – 2/3 oz. for limited control of certain small grass weeds. See Table 2, under the section ADDITIONAL RECOMMENDATIONS for details.

As weeds mature, their sensitivity to HELM NICOSULFURON 75 decreases. As grassy weeds become mature (more than 3 tillers), they may not reach the size listed below, due to drought or other environmental factors. Grassy weeds that are maturing rapidly should be treated before they reach the stages listed below.

When applied as directed, HELM NICOSULFURON 75 will control the following weeds:

**Table 1. Weeds controlled with 2/3 ounces HELM Nicosulfuron 75**

Grasses	Maximum Height
Barnyardgrass	4"
Broadleaf signalgrass	2"
Foxtails (bristly, giant, green, yellow)	4"

(continued)

**Table 1. Weeds controlled with 2/3 ounces HELM Nicosulfuron 75 (continued)**

Grasses	Maximum Height
Itchgrass	6"
Johnsongrass seedling	12"
rhizome	18"
Panicum (Texas, browntop)	3"
fall	4"
Quackgrass*	10"
Ryegrass (Italian, perennial)	6"
Sandbur (field, longspine)*	3"
Shattercane	12"
Sorghum alnum	12"
Timothy	6"
Volunteer cereals (wheat, barley, oats)	6***
Wild oats	4"
Wild proso millet	4"
Wirestem muhly*	8"
Witchgrass	6"
Woolly Cupgrass*†	4"

\* Requires the use of COC plus ammonium nitrogen fertilizer. Cultivation or re-treatment may be required. See “FOR ADDITIONAL CONTROL OF LATER EMERGING GRASSES”

\*\*10 inches in the states of WA, OR, ID, and MT, where the use of MSO adjuvants are preferred. See SPRAY ADJUVANTS.

Broadleaves	Maximum Height or Diameter
Burcucumber	3"
Dandelion	6"
Hemp dogbane*	4"
Jimsonweed	3"
Morningglory (ivyleaf, pitted)	3"
Tall	2"
Pigweed (redroot, smooth)	4"
Pokeweed*	4"
Smartweeds (ladysthumb, PA)	4"
Thistle, Canada*	4"

\*Suppression

#### **Popcorn, Field Corn Grown for Seed and Sweet Corn**

HELM NICOSULFURON 75 may be broadcast or applied with drop nozzles to popcorn or field corn grown for seed that is less than 20" tall (free-standing) or that exhibits up to and including 5 leaf-collars (V5), whichever is most restrictive. Do not apply to corn that is taller than 20" or that exhibits more than 5 leaf-collars (V5), whichever is more restrictive.

Many seed companies have tested seed corn inbreds or yellow popcorn hybrids for sensitivity to HELM NICOSULFURON 75 and have reported excellent safety. Do not apply HELM NICOSULFURON 75 to any white popcorn inbred, or white popcorn hybrid unless specifically approved by the seed company. This includes "White Dynamite" popcorn.

HELM NICOSULFURON 75 may be applied to certain sweet corn hybrids grown for fresh markets or under contract for processing. Applications of HELM NICOSULFURON 75 may be applied broadcast or with drop nozzles (post-directed) on sweet corn up to 12 inches tall or up to and including 5 leaf-collars (V5). For sweet corn 12-18 inches tall, apply only with drop nozzles. Do not apply to sweet corn taller than 18 inches or those which exhibit 6 or more leaf-collars (V6), and make only one application of HELM NICOSULFURON 75 per year.

Sweet corn hybrid sensitivity to HELM NICOSULFURON 75 is highly variable, and not all hybrids have been tested for crop tolerance.

Not all seed corn inbreds, popcorn, or sweet corn hybrids have been tested, nor does Sharda USA have access to all seed company data. Consequently, Sharda USA is not responsible for any crop injury arising from the use of HELM Nicosulfuron 75 on field corn grown for seed, popcorn, or sweet corn. When tank mixing, check the tank mix partner label for tolerances and instructions for use.

See SOIL INSECTICIDE INTERACTION INFORMATION regarding the use of HELM Nicosulfuron 75 on popcorn, sweet corn or field corn grown for seed that has been previously treated with a soil insecticide.

#### **SPRAY ADJUVANTS**

Applications of HELM Nicosulfuron 75 must include either a crop oil concentrate or a nonionic surfactant. In addition, an ammonium nitrogen fertilizer must be used unless specifically prohibited by tank mix partner labeling. Crop oil concentrate plus ammonium nitrogen fertilizer is the preferred adjuvant system for activity on difficult to control species such as woolly cupgrass, quackgrass, sandbur and wirestem muhly. If another herbicide is tank mixed with HELM Nicosulfuron 75, select adjuvants authorized for use with both products. Products must contain only EPA-exempt ingredients (40 CFR 1001).

***Crop Oil Concentrate (COC) - Petroleum or Modified Seed Oil (MSO)***

- Apply at 1% v/v (1 gallon per 100 gallons spray solution) or 2% under arid conditions.
- MSO adjuvants may be used at 0.5% v/v (0.5 gallons per 100 gallons spray solution) if specifically noted on adjuvant product labeling.
- Oil adjuvants must contain at least 80% high quality, petroleum (mineral) or modified vegetable seed oil with at least 15% surfactant emulsifiers.

***Nonionic Surfactant (NIS)***

- Apply at 0.25% v/v (1 quart per 100 gallons spray solution) or 0.5% under arid conditions.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

***Ammonium Nitrogen Fertilizer***

- Use 2 quarts/acre of a high-quality urea ammonium nitrate (UAN), such as 28%N or 32%N, or 2 pounds/acre of a spray-grade ammonium sulfate (AMS). Use 4 quarts/acre UAN or 4 pounds/acre AMS under arid conditions.
- Do not use liquid nitrogen fertilizer as the total carrier solution.

***Special Adjuvant Types***

- Combination adjuvant products may be used at doses that provide the required amount of NIS, COC, MSO and/or ammonium nitrogen fertilizer. Consult product literature for use rates and restrictions.

***MIXING INSTRUCTIONS***

1. Fill the tank 1/4 to 1/3 full of water.
2. While agitating, add the required amount of HELM Nicosulfuron 75.
3. Continue agitation until HELM Nicosulfuron 75 is fully dispersed, at least 5 minutes.

4. Once HELM Nicosulfuron 75 is fully dispersed, maintain agitation and continue filling tank with water. Thoroughly mix HELM Nicosulfuron 75 with water before adding any other material.
5. As the tank is filling, add the required spray adjuvants (crop oil concentrate, non-ionic surfactant, or ammonium nitrogen fertilizer).
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply HELM Nicosulfuron 75 spray mixture within 24 hours of mixing to avoid product degradation.
8. If HELM Nicosulfuron 75 and a tank mix partner are to be applied in multiple loads, pre-slurry HELM Nicosulfuron 75 in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of HELM Nicosulfuron 75.

***SEQUENTIAL APPLICATIONS FOLLOWING REDUCED RATES OF PREEMERGENCE HERBICIDES***

HELM Nicosulfuron 75 may be used as a sequential application in a planned post-emergence weed control program in corn following a reduced rate of a preemergence herbicide.

Apply a reduced rate of a preemergence grass herbicide prior to corn emergence and then follow with a postemergence application of HELM Nicosulfuron 75. Apply products such as "Cinch"®, "Cinch"® ATZ, "Balance" PRO, "Axiom", "Dual" H Magnum, "Surpass", "Outlook" and "Harness" Xtra at as low as 1/4 to 1/2 of the full labeled use rate and follow with a sequential postemergence application of HELM Nicosulfuron 75. Refer to the preemergence grass herbicide label for use restrictions, application information, rotational crop guidelines, and cautionary statements prior to applying HELM Nicosulfuron 75.

Do not apply HELM Nicosulfuron 75 to corn that exhibits herbicide injury from previous applications made to the current or preceding crop.

**TANK MIX APPLICATIONS*****For Additional Control of Broadleaf Weeds***

Nicosulfuron WDG may be tank mixed with many herbicides registered for post-emergence application in corn for additional control of broadleaf weeds. See the tank mix partner label for weeds controlled, precautions, use restrictions, adjuvant and crop rotation information. The most restrictive language on either label shall apply.

In addition to the tank mixtures noted above, HELM Nicosulfuron 75 may be tank mixed with the rates of products listed below for improved control of many broadleaf weeds, including cocklebur, dandelion, Eastern black nightshade, lambsquarters, pigweeds, ragweeds, PA smartweed and velvetleaf. See ADDITIONAL RECOMMENDATIONS AND/OR RECOMMENDATIONS FOR SPECIFIC WEED PROBLEMS below for additional information.

Product	Rate/A
Atrazine*	Up to 2 lb. a.i.
dicamba (e.g., "Clarity" - 4 lb/gal dicamba)	2-4 fl. oz.
dicamba + atrazine (e.g. "Marksman"-1.1 lb/gal dicamba)*	8-16 fl. oz.
"Callisto"	1.5-3.0 fl. oz.
"Distinct"***	1-2 oz.
"Exceed"***	0.25-0.5 oz.
"Northstar"***	2.5-5.0 oz.

\*Make applications to emerged corn before the corn reaches 12" tall.

\*\*Do not apply to sweet corn, seed corn, or popcorn.

Rates listed are for the specific products noted in the table. If other brands or formulations are used, rates of active ingredients should be adjusted to correspond to the products indicated. Formulations of products other than those listed may not have

been tested with HELM Nicosulfuron 75. Check with the manufacturer for information on tank mix compatibility prior to using (see TANK MIX COMPATIBILITY TESTING).

Crop oil concentrate plus ammonium nitrogen fertilizer is the preferred adjuvant for tank mixtures when using products at the low end of the rate range indicated in the table. The use of nonionic surfactant is permitted in place of crop oil concentrate for tank mixtures containing dicamba, however, overall weed control may be reduced. See SPRAY ADJUVANTS for adjuvant rate recommendation.

Do not use MSO adjuvants when tank mixing HELM Nicosulfuron 75 with >1.5 ounces "Callisto".

**RECOMMENDATIONS FOR SPECIFIC WEED PROBLEMS*****Reduced Rates of HELM Nicosulfuron 75***

HELM Nicosulfuron 75 may be applied at 1/3 - 2/3 ounces for control of the small grass weeds noted in the table below. Always use a crop oil concentrate plus ammonium nitrogen fertilizer when applying reduced rates of HELM Nicosulfuron 75.

**Table 2. Weeds controlled with reduced rates of HELM Nicosulfuron 75**

	Maximum Height or Diameter HELM Nicosulfuron 75 Rate		
	1/3 oz.	1/2 oz.	2/3 oz.
<b>Grasses</b>			
Barnyardgrass	2"	3"	4"
Foxtails (bristly, giant)	2"	3"	4"
yellow	—	2"	4"
Itchgrass	2"	4"	6"
Johnsongrass, seedling	—	8"	12"
rhizome		8"	18"

(continued)

**Table 2. Weeds controlled with reduced rates of HELM Nicosulfuron 75 (continued)**

	Maximum Height or Diameter HELM Nicosulfuron 75 Rate		
	1/3 oz.	1/2 oz.	2/3 oz.
<b>Grasses</b>			
Panicum , Texas fell	1"	2"	3"
Sandbur ,field	—	1"	3"
Shattercane	3"	6"	12"
Sorghum alnum	3"	6"	12"
Timothy	2"	4"	6"
Volunteer cereals	—	2"	6"
Wild oats	2"	3"	4"
Wild proso millet	-	2"	4"
Witchgrass	2"	4"	6"
Woolly cupgrass	-	-	4"

***Tank Mixtures with Atrazine***

HELM Nicosulfuron 75 may be tank mixed with up to 2 pounds a.i. atrazine\* for additional control of many broadleaf weeds, including:

	Maximum Height or Diameter
<b>Sicklepod</b>	2"
<b>Prickly sida</b>	2"
<b>Wild Radish</b>	12"
<b>Cutleaf evening</b>	6"
<b>Florida pusley</b>	2"

\*For best results add 0.25 - 2.0 quarts Atrazine 4L OR 4 - 35 ounces Atrazine 90DF. Products containing atrazine are restricted use products.

HELM Nicosulfuron 75 + atrazine tank mixtures may result in reduced control of grasses (antagonism) if applied to grasses under low moisture stress or to grasses exceeding the maximum labeled height. Before applying Nicosulfuron + atrazine tank mixtures, refer to the atrazine product label for information regarding the maximum amount of atrazine that may be applied in a season.

***Tank Mixtures with "Callisto"***

HELM Nicosulfuron 75 may be tank mixed with 1.5 - 3.0 fluid ounces/acre of "Callisto" herbicide for weed control as indicated in the table below:

Species	Maximum Height or Diameter					
	"Callisto" Alone			"Callisto" + atrazine*		
	1.5 oz	2.0 oz	3.0 oz	1.5 oz	2.0 oz	3.0 oz
Cocklebur	4"	4"	4"	10"	10"	10"
Dandelion	10"	10"	10"	10"	10"	10"
Jimsonweed	4"	4"	4"	4"	10"	10"
Kochia	-	-	4"	-	4"	4"
Lambsquarters, common	4"	4"	4"	10"	10"	10"
Morningglory ivyleaf	4"	4"	4"	4"	4"	4"
Mustard, wild	-	-	4"	-	-	10"

*(continued)*

Species	Maximum Height or Diameter					
	"Callisto" Alone			"Callisto" + atrazine*		
	1.5 oz	2.0 oz	3.0 oz	1.5 oz	2.0 oz	3.0 oz
Nightshade, black, Eastern black	4"	4"	4"	10"	10"	10"
Pigweed, palmer	-	-	4"	4"	4"	4"
Pigweed, redroot, smooth	4"	4"	4"	10"	10"	10"
Ragweed, common	-	-	-	4"	10"	10"
Ragweed, giant	-	3"	4"	4"	10"	10"
Smartweed, ladythumb	-	4"	4"	4"	10"	10"
Smartweed, Pennsylvania	4"	4"	4"	4"	10"	10"
Sunflower, common	4"	4"	4"	4"	4"	10"
Velvetleaf	4"	4"	4"	10"	10"	10"
Waterhemp, tall & common		4"	4"	4"	10"	10"

\*Plus 0.25 to 0.75 pounds a.i. atrazine per acre, may provide better control when weeds are at maximum height.

For improved grass and broadleaf weed control, HELM Nicosulfuron 75 tank mixtures with 1.5 ounces "Callisto" (with or without atrazine) may be applied with 0.5% v/v MSO spray adjuvant. Do not use MSO adjuvants when tank mixing HELM Nicosulfuron 75 with >1.5 ounces "Callisto". Use a petroleum-based crop oil concentration + an ammonium nitrogen fertilizer.

#### **Tank Mixes with "Impact" plus atrazine**

HELM Nicosulfuron 75 may be tank mixed with 0.5 to 0.75 fl. oz./acre of "Impact" herbicide plus atrazine at 0.375 to 1.5 lbs. a.i./acre for weed control as indicated in the table below:

Species	HELM Nicosulfuron 75 + Atrazine +	
	"Impact" 0.5 oz.	"Impact" 0.75 oz.**
Amaranth, Palmer	4"	6"
Cocklebur, common	5"	8"
Jimsonweed	4"	6"
Kochia	4"	6"
Lambsquarter, common	4"	6"
Morningglory, annual	4"	4"
Mustard, wild	4"	6"
Nightshade (black, Eastern black)	4"	6"
Pigweed (redroot, smooth)	4"	6"
Ragweed, common	4"	6"
Ragweed, giant	5"	8"
Smartweed, Pennsylvania	2"	3"

(continued)

Species	HELM Nicosulfuron 75 + Atrazine +	
	"Impact" 0.5 oz.	"Impact" 0.75 oz.**
Smartweed, Ladysthumb	2**	3"
Sunflower, common	5**	8"
Thistle, Canada	4**	6"
Velvetleaf	5"	8"
Waterhemp (tall, common)	4"	6"

'suppression

\*Refer to Impact label for additional information regarding tank mixtures, adjuvants, and rotational crops. Current research supports applications at these use rates only within the following geographies: Illinois, north of I-80; Iowa, north of I-80 (excluding the area that is both north of U.S. Hwy. 20 and west of U.S. Hwy. 71); Michigan, entire state; Minnesota, east of U.S. Hwy. 71; Nebraska, north of Hwy. 92; Wisconsin, entire state.

\*\*Refer to Impact herbicide label for specific rotational crop information.

#### **Tank Mixtures with "Lumax" or "Lexar"**

HELM Nicosulfuron 75 may be tank mixed with 2 pints/acre of "Lumax" or 2 1/3 pints/acre of "Lexar" herbicide for weed control as indicated in the table below:

Species	"Lumax" 2 pints	"Lexar" 2 1/3 pints
Amaranth, Palmer	4"	4"
Cocklebur, common	10"	10"
Dandelion	10"	10"
Jimsonweed	10"	10"
Kochia	4**	4"

(continued)

Species	"Lumax" 2 pints	"Lexar" 2 1/3 pints
Lambsquarter, common	10"	10"
Morningglory, annual	4"	4"
Mustard, wild	4"	10"
Nightshade (black, Eastern black)	10"	10"
Pigweed (redroot, smooth)	10"	10"
Ragweed, common	10"	10"
Ragweed, giant	10"	10"
Smartweed, Pennsylvania	10"	10"
Smartweed, Ladysthumb	10"	10"
Sunflower, common	4"	4"
Velvetleaf	10"	10"
Waterhemp (tall, common)	10"	10"

#### **For Additional Control of Later Emerging Grasses**

HELM Nicosulfuron 75 may be tank mixed with full or reduced rates of preemergence grass herbicides labeled for early postemergence application to field corn (such as Cinch®, Cinch® ATZ, "Prowl", "Surpass" EC, "Dual" II Magnum, or "Outlook") for residual activity on later-emerging flushes of grass. Application must be made before the grass emerges and before other grass weeds on the HELM Nicosulfuron 75 label exceed their labeled sizes.

The use of nonionic surfactant is recommended in place of crop oil concentrate for tank mixtures with preemergence grass herbicides where applications are made early postemergence to small grass weeds. See SPRAY ADJUVANTS for adjuvant rate recommendations.

When tank mixing HELM Nicosulfuron 75 with EC formulated preemergence grass herbicides such as Cinch®, "Prowl", or "Dual" II Magnum, do not add "Callisto" herbicide to the tank mixture. When other formulations of preemergence grass herbicides are tank mixed with HELM Nicosulfuron 75 + "Callisto" (such as Cinch® ATZ or "Bicep II Magnum"), limit preemergence herbicide rates to 2/3 times full rates, always add nonionic surfactant in place of crop oil concentrate, and, limit broadleaf weed sizes to less than or equal to 4" tall.

When tank mixing HELM Nicosulfuron 75 with "Lumax" or "Lexar" herbicides, limit "Lumax" rates to no more than 2 pints and "Lexar" rates to no more than 2 1/3 pints/acre. Always add nonionic surfactant in place of crop oil concentrate, omit adjuvants containing ammonium nitrogen fertilizer, and limit applications to corn up to 5" tall.

Tank mixes of HELM Nicosulfuron 75 and preemergence grass herbicides must be broadcast applied postemergence to field corn before the crop exceeds the heights listed on the preemergence grass herbicide label. Refer to WHEN TO APPLY-POSTEMERGENCE and the preemergence grass herbicide label for complete postemergence application information, rates, and restrictions.

#### **Tank Mixtures with Insecticides**

HELM Nicosulfuron 75 may be tank mixed with pyrethroid or carbamate insecticides. See Soil Insecticide Interaction section for information on use of HELM Nicosulfuron 75® following soil insecticides application.

#### **Other Tank Mixtures**

Other than the exceptions noted, and in addition to the tank mix partners and rates indicated above, Nicosulfuron 75 WDG may be tank mixed or followed with sequential applications of other products registered for use in field corn. Applications of full or reduced rates of other products registered for use in corn provided:

- The tank mix product is labeled for the same timing, method of application, adjuvants, and use restrictions as HELM Nicosulfuron 75.
- The tank mixture is not specifically prohibited on the label of the tank mix product.
- The tank mix combination is compatible as determined by a "jar test" described in the TANK MIX COMPATIBILITY TESTING section below.

Weed control and crop response with tank mixtures not specifically recommended in this label are the responsibility of the user and manufacturer of the tank mix product.

#### **Tank Mixing Precautions**

A corn plant's predisposition to develop fused tissue emerging from the whorl (rattail) after the V-11 stage may increase when a product containing dicamba (i.e. "Clarity", "Marksman") is applied to small corn under early stressful conditions. Be aware of this when applying tank mixes with dicamba to small corn (V-3 stage or smaller) under stressful conditions. See ENVIRONMENTAL CONDITIONS for a description of these stressful conditions.

To avoid crop injury or antagonism, apply the products indicated below at least seven days before or three days after the application of HELM Nicosulfuron 75.

- Do not tank mix HELM Nicosulfuron 75 with "Basagran" and "Laddock" or severe crop injury may occur.
- Do not tank mix HELM Nicosulfuron 75 with 2,4-D- containing products as severe grass control antagonism may occur.
- Do not tank mix HELM Nicosulfuron 75 with foliar-applied organophosphate insecticides such as "Lorsban", malathion, parathion, etc., as severe crop injury may occur.

Do not exceed labeled application rates. Do not tank mix HELM Nicosulfuron 75 with other products that contain the same active ingredient as HELM Nicosulfuron 75 (nicosulfuron) unless the label of either tank mix partner specifies the maximum rate that may be used.

**TANK MIX COMPATIBILITY TESTING**

Perform a jar test prior to tank mixing to ensure compatibility of HELM Nicosulfuron 75 and other pesticides. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

**SEQUENTIAL HELM NICOSULFURON 75 APPLICATIONS**

Annual grasses may have more than one flush of emerging seedlings. Also, regrowth of treated annual grasses may occur due to adverse environmental conditions following application. Perennial grasses may regrow from underground stems or roots, depending upon environmental conditions. To control grasses under these conditions, a sequential application of HELM Nicosulfuron 75 may be necessary. The combined dosage of the sequential applications cannot exceed 1 1/3 ounces per acre of HELM Nicosulfuron 75.

**CULTIVATION**

A timely cultivation may be necessary to control suppressed weeds, or weeds that emerge after an application of HELM Nicosulfuron 75.

Optimum timing for cultivation is 7-14 days after HELM Nicosulfuron 75 application or upon seeing the establishment of new weeds.

**ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY**

HELM Nicosulfuron 75 provides best results when applied to young, actively growing weeds. Applications made during warm, moist conditions (70°F or more) and adequate soil moisture both before and after application maximizes performance.

The degree and duration of control depend on spray coverage, weed spectrum, weed size, growing conditions before and after treatment, soil moisture, and adjuvant selection.

HELM Nicosulfuron 75 is rainfast in 4 hours.

Treating weeds that exceed maximum label height or that are under stress may result in incomplete control. Poor weed control or crop injury may result from applications made to plants under stress from:

- abnormally hot or cold weather
- environmental conditions such as drought, water-saturated soils, hail damage, or frost
- disease, insect, or nematode injury
- prior herbicide, or carryover from a previous year's herbicide application

Severe stress from conditions preceding or immediately following application may also result in crop injury or poor weed control. Stress affects all weeds, but especially weeds such as woolly cupgrass, green and yellow foxtail, and wild proso millet.

If the corn or grass weeds are under stress, delay application until stress passes and both weeds and corn resume active growth.

HELM Nicosulfuron 75 rapidly inhibits the growth of susceptible weeds, reducing weed competition within as little as 6 hours after application. Susceptible plants are controlled in 7-21 days.

**SOIL INSECTICIDE INTERACTION INFORMATION**

Before using HELM Nicosulfuron 75, ensure that it is compatible with any insecticides previously applied to the corn crop.

HELM Nicosulfuron 75 may interact with certain insecticides previously applied to the crop. Crop response varies with field corn type, insecticide used, insecticide application method, and soil type.

HELM Nicosulfuron 75 may be applied to corn previously treated with "Fortress", "Aztec", or "Force" insecticides or nonorganophosphate (OP) soil insecticides regardless of soil type.

- DO NOT APPLY HELM Nicosulfuron 75 to corn previously treated with "Counter" 15G or to corn treated with "Counter" 20CR in-furrow or over the row at cultivation.
- Applications of HELM Nicosulfuron 75 to corn previously treated with "Counter" 20 CR, "Lorsban", or "Thimet" may cause unacceptable crop injury, especially on soils of less than 4% organic matter.

#### **CROP ROTATION**

Rotational crops vary in their response to low concentrations of HELM Nicosulfuron 75 remaining in the soil. HELM Nicosulfuron 75 dissipates rapidly in warm, acidic, microbiologically active soils.

The amount of HELM Nicosulfuron 75 which may be present in the soil depends on application rate, soil pH and organic matter content, elapsed time since application, crop production practices, and environmental factors.

Injury to rotational crops may occur in high-pH, cold soils if dry weather prevails between application and rotational crop planting.

Soil pH should be determined by laboratory analysis using the 1:1 soil: water suspension method on representative soil samples taken at 0-4" depth. Soil pH varies within fields; therefore, recropping should be based on the highest soil pH within each field. Consult local extension publications for recommended soil sampling procedures.

The following rotational intervals should be observed when using HELM Nicosulfuron 75 at a maximum of 1 1/3 ounces:

#### **HELM NICOSULFURON 75 ROTATIONAL CROP GUIDELINE- A No Soil pH Restrictions**

<b>Crop Rotational</b>	<b>Interval in Months</b>
Corn (field, seed)	Anytime
Corn (pop, sweet)*	10
Soybeans	0.5 (15 days)
Cereals, spring (barley, oats, rye)	8
Cereals, winter (barley, oats, rye)	4
Cotton	10
Dry Beans, Peas, Snap Beans	10 12
Red Clover**	12
Other Crops	See Rotational Crop Guidelines B and C

\* Except the sweet corn varieties "Merit", "Carnival", and "Sweet Success", for which the minimum time interval is 15 months.

\*\*Except for the state of Kansas east of Highway 75, for Minnesota east and south of the Red River Valley and for the states east of the line formed by the western borders of Iowa, Missouri, Arkansas, and Louisiana, where the minimum time interval is 10 months.

**HELM NICOSULFURON 75 ROTATIONAL CROP GUIDELINE – B**  
**With Soil pH  $\leq$  7.5 Restrictions**

Crop	Rotational Interval in Months	
	pH 7.5	pH >7.5
Sorghum	10	18*
Sunflowers	11**	18
All other crops not listed in Rotational Guidelines A or B	See Rotational Guideline C	

\*Except in Texas and Oklahoma east of Highway 281, where the rotational interval is 10 months, regardless of pH.

\*\*Precipitation following application must exceed 14" prior to planting sunflowers.

**HELM NICOSULFURON 75 ROTATIONAL CROP GUIDELINE – C**  
**With Soil pH  $\leq$  6.5 Restrictions**

Crop	Rotational Interval in Months	
	pH 6.5	pH >6.5
Sugarbeets*, Potatoes**	10	18***
All other crops not listed in Rotational Guidelines A or B	10	18

\*Except on irrigated sites in Colorado, Wyoming, Nebraska, Texas, Michigan, and Ohio, where precipitation following application must exceed 25" prior to planting beets, where the interval is 10 months on soils with pH < 7.5. Sites in Minnesota east and south of the Red River Valley may follow these guidelines provided maximum rates of HELM Nicosulfuron 75 do not exceed 0.67 oz.

\*\*Irrigated potatoes following irrigated corn treated in the States of WA, OR, ID, or Utah can be planted 10 months after using HELM Nicosulfuron 75 on sprinkler irrigated corn with no soil pH restrictions, providing the maximum use rate on corn does not exceed 1.0 ounce product per season. Corn treated with HELM Nicosulfuron 75 must be grown to maturity and receive a minimum of 18 inches of irrigation water before potatoes can be planted at this rotation interval. Injury to potatoes may occur if less than 18 inches of irrigation is used on the previous corn crop. HELM Nicosulfuron 75 may not be used in a tankmix or sequential application program with other ALS-inhibiting herbicides such as "Exceed" or "Beacon".

\*\*\*In North Dakota and northwest Minnesota, the cumulative precipitation in the 18 months following application must exceed 28" in order to rotate to sugarbeets or potatoes.

**ROTATIONAL CROP GUIDELINES-D**  
**Maximum Use Rate: 0.67 ounces**

These guidelines may be observed when using a single application of HELM Nicosulfuron 75 per cropping season with a maximum use rate of 0.67 ounces product. Rotational intervals should be extended to 12 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

Crop	Rotational Interval in Months
Alfalfa*	10
Canola	10
Flax**	10
Potato	10
Red clover	10
Sunflower	10

\*On sprinkler irrigated fields in Idaho, Utah, and Northern Nevada it is best to use deep fall tillage such as plowing prior to planting alfalfa. Product degradation may be less on furrow irrigated soils and may result in some crop injury.

\*\*Rotational intervals should be extended to 18 months if drought conditions prevail after application and before the rotational crop is planted, unless sprinkler irrigation has been applied and totals greater than 15" during the growing season.

#### **APPLICATION INFORMATION**

Many crops are highly sensitive to HELM Nicosulfuron 75. All direct or indirect contact (such as spray drift) with crops other than field corn should be avoided (see also SPRAY DRIFT MANAGEMENT).

For all application systems, use 50-mesh or larger strainer screens.

Do not apply HELM Nicosulfuron 75 through any type of irrigation system.

#### **GROUND APPLICATION**

##### **Broadcast Application**

- Use a minimum of 15 gallons of water per acre (15 GPA) for best performance. Use a minimum of 10 gallons of water per acre (GPA) for light, scattered stands of weeds.
- For best performance, select nozzles and pressure that deliver MEDIUM spray droplets, for example, as indicated in nozzle manufacturer's catalogues and in accordance with AS AE Standard S572. Nozzles that deliver COARSE spray droplets may be used to reduce drift, provided spray volume is increased to maintain coverage on small weeds. For optimal product performance and minimal spray drift, adjust the spray boom to the lowest possible spray height recommended in manufacturers' specifications.
- Ensure that equipment is set up to avoid applying an excessive rate directly over the rows and into the corn plant whorl. This is most likely to occur when a nozzle is positioned directly above the row.

- Overlaps or starting, stopping, slowing, and turning while spraying may result in crop injury.

##### **Band Application**

For band applications, use proportionately less spray mixture, and carefully calibrate the band applicator to not exceed the labeled rate. Carefully follow the manufacturer's instructions for nozzle type (flat fans), orientation, distance of nozzles from the crop and weeds, spray volumes, calibration and spray pressure.

##### **AERIAL APPLICATION**

In New York and California aerial application is not permitted.

Use nozzle types and arrangements that will provide optimum spray distribution and maximum coverage at a minimum of 3 GPA.

Do not apply during a temperature inversion, when winds are gusty, or when conditions favor poor coverage and/or off-target spray movement.

##### **SPRAYER PREPARATION/CLEANUP**

It is important that spray equipment is clean and free of previous pesticide deposits before using HELM Nicosulfuron 75 and then properly cleaned out following application. Clean all application equipment before applying HELM Nicosulfuron 75. Follow the cleanup procedures specified on the label of the product previously sprayed. If no cleanup procedure is provided, use the procedure that follows. Immediately following applications of HELM Nicosulfuron 75, thoroughly clean all mixing and spray equipment to avoid subsequent crop injury.

##### **Note:**

- When cleaning spray equipment before applying HELM Nicosulfuron 75, read and follow label directions for proper rinsate disposal of the product previously sprayed.
- Steam cleaning of aerial spray tanks will help to dislodge any visible pesticide deposits.

- When spraying or mixing equipment will be used over an extended period to apply multiple loads of HELM Nicosulfuron 75, partially fill the tank with fresh water at the end of each day of spraying, flush the boom and hoses, and allow to sit overnight.

#### **Cleanup Procedure**

1. Drain the tank and thoroughly hose down the interior surfaces. Flush the tank, hoses, and boom with clean water for a minimum of 5 min.
2. Partially fill the tank with clean water and add one gallon of household ammonia\* (containing 3% active) for every 100 gallons of water. Finish filling the tank with water, then flush the cleaning solution through the hoses, boom, and nozzles. Add more water to completely fill the tank and allow to agitate/recirculate for at least 15 min. Again, flush the hoses, boom, and nozzles with the cleaning solution, then drain the tank.
3. Repeat Step 2.
4. Remove the nozzles and screens and clean separately in a bucket containing the cleaning agent and water.
5. Thoroughly rinse the tank with clean water for a minimum of 5 min, flushing the water through the hoses and boom.

\* Equivalent amounts of an alternate strength ammonia solution may be used.

#### **SPRAY DRIFT MANAGEMENT**

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

#### **IMPORTANCE OF DROPLET SIZE**

The most effective way to reduce drift potential is to apply large droplets (>150 - 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the

environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See **Wind, Temperature and Humidity**, and **Temperature Inversions** sections of this label.

#### **Controlling Droplet Size - General Techniques**

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

#### **Controlling Droplet Size - Aircraft**

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- **Application Height** - Application more than 10 ft above the canopy increases the potential for spray drift.

**BOOM HEIGHT**

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

**WIND**

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

**TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

**TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning.

Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

**INTEGRATED PEST MANAGEMENT**

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

**RESISTANCE**

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

#### **IMPORTANT PRECAUTIONS**

Injury to or loss of desirable vegetation may result from failure to observe the following:

- Do not apply HELM Nicosulfuron 75 or drain or flush application equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, tennis courts, or similar areas.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water.
- Thoroughly clean application equipment immediately after use. (See the Sprayer Cleanup section of this label for instructions).
- Do not graze or feed forage, hay, or straw from treated areas to livestock within 30 days of HELM Nicosulfuron 75 application.

## **STORAGE AND DISPOSAL**

Do not contaminate water, other pesticides, fertilizer, food or feed in storage.

**Pesticide Storage:** Store product in original container only. Store in a cool, dry place.

**Pesticide Disposal:** Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

*(continued)*

## **STORAGE AND DISPOSAL** *(continued)*

#### **Container Disposal:**

**For Plastic Containers:** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Dispose of bags at an approved waste disposal facility, in accordance with Federal, state and local regulations.

**For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

**For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above.

**For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Cinch®, Asana® and Lannate® are trademarks or registered trademarks of E. I. duPont de Nemours and Company.

"Axiom", "Aztec", "Balance" are trademarks or registered trademarks of Bayer CropScience.

"Basagran", "Clarity", "Counter", "Distinct", "Laddok", "Marksman", "Outlook", "Prowl", "Thimet" are trademarks or registered trademarks of BASF Ag Products.

"Beacon", "Callisto", "Dual", "Exceed", "Force", "Lumax", "Lexar", and "Northstar" are trademarks or registered trademarks of Syngenta Crop Protection Inc.

"Fortress" and "Impact" are registered trademarks of Amvac Chemical Corporation.

"Harness" is a registered trademark of Monsanto Technology LLC.

"Lorsban", "Surpass" are registered trademarks of Dow AgroSciences LLC.

### **LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off target movement, unconventional farming techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Helm Agro US, Inc.. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants. Helm Agro US, Inc. does not agree to be an insurer of these risks. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

Helm Agro US, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, HELM AGRO US, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL SHARDA USA OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF SHARDA USA OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF HELM AGRO US, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

**notes**

# HELM Nicosulfuron 75

Water Dispersible Granule Herbicide for  
Use on Corn

**ACTIVE INGREDIENT:**

Nicosulfuron 2-[[[4,6-dimethoxyimidin-2-yl)aminocarbonyl]aminosulfonyl] N,N-dimethyl-3-pyridinecarboxamide ..... 75%

**INERT INGREDIENTS:** ..... 25%

**TOTAL** .....100%

**STOP - Read the label before use**  
**KEEP OUT OF REACH**  
**OF CHILDREN**  
**CAUTION**

EPA Reg. No. 74530-26 EPA Est. No. 083390-CHN-001

**Net Contents: 10 ou.**



**Distributed by:**  
Helm Agro US, Inc.  
8295 Tournament Dr. Suite #310  
Memphis, TN 38125

Manufactured for:

**Sharda USA LLC**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
Caution! Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

**FIRST AID**

**If In Eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **Have a product container or label with you when calling a poison control center or doctor, or going for treatment. Note To Physician: No specific antidote is available. Treat the patient symptomatically.**

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater. Do not apply where/when conditions could favor runoff.

**DIRECTIONS FOR USE**

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

## STORAGE AND DISPOSAL

Do not contaminate water, other pesticides, fertilizer, food or feed in storage. **Pesticide Storage:** Store product in original container only. Store in a cool, dry place. **Pesticide Disposal:** Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal: For Plastic Containers:** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **For Fiber Sacks:** Completely empty fiber sack by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Dispose of bags at an approved waste disposal facility, in accordance with Federal, state and local regulations. **For Fiber Drums With Liners:** Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by state and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner. **For Bags Containing Water Soluble Packets:** Do not reuse the outer box or the resealable plastic bag. When all water-soluble packets are used, the outer packaging should be clean and may be disposed of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by open burning. If burned, stay out of smoke. If the resealable plastic bag contacts the formulated product in any way, the bag must be triple-rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer wrap as described above. **For Metal Containers (non aerosol):** Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

OPEN HERE  
PRESS TO RESEAL